


Series 2, Mar 27th, 2020 (Kernel)

Problem 1 (SVM):

This exercise is based on an exercise designed by Stephanie Hyland. In its original formulation, the perceptron aims to minimise a 0/1-loss function (shown below, solid). Because this objective is neither convex nor differentiable, a surrogate loss function is optimised (typically, $l_p(\mathbf{w}; \mathbf{x}, y) = \max(0, -y\mathbf{w}^T \mathbf{x})$, dashed). In this exercise, we consider a different surrogate loss function l_s , which approximates the 0/1-loss function more closely.

$$l_s(\mathbf{w}; \mathbf{x}, y) = \begin{cases} 0, & \text{for } \text{sign}(\mathbf{w}^T \mathbf{x}) = y \\ \sqrt{-y\mathbf{w}^T \mathbf{x}}, & \text{for } \text{sign}(\mathbf{w}^T \mathbf{x}) \neq y \end{cases}$$



1. Mark the following statements as True or False. Try to justify the answer for yourself.

- (a) l_p is convex.
- (b) l_p is differentiable.
- (c) l_s is convex.
- (d) l_s is differentiable.

Solution:

Only (a) is True.

- (a) l_p , known as *the hinge loss*, is convex because it is the maximum of two linear functions, and:
 - i. Any linear function is convex.
 - ii. The maximum of two convex functions is convex.
- (b) Let's differentiate with respect to $y\mathbf{w}^T\mathbf{x}$. If $\text{sign}(\mathbf{w}^T\mathbf{x}) = y$, $l'_p(\mathbf{w}; \mathbf{x}, y) = 0$. If $\text{sign}(\mathbf{w}^T\mathbf{x}) \neq y$, $l'_p(\mathbf{w}; \mathbf{x}, y) = -1$.
To check differentiability, we need to check the limit at point 0. $\lim_{x \rightarrow 0^-} l'_p = -1$. l_p is not differentiable at $y\mathbf{w}^T\mathbf{x} = 0$, since the left and right derivatives are not equal.
- (c) l_s is not convex.
To check whether l_s is convex, we can look at $f(x) = \sqrt{x}$.
A way to show that $f(x) = \sqrt{x}$ is not convex is to show that $-f(x)$ is convex.

$$\begin{aligned} \sqrt{tx_1 + (1-t)x_2} &> t\sqrt{x_1} + (1-t)\sqrt{x_2} \\ tx_1 + (1-t)x_2 &> t^2x_1 + (1-t)^2x_2 + t1(1-t)\sqrt{x_1x_2} \\ x_1 + x_2 &> 2\sqrt{x_1x_2} \\ (\sqrt{x_1} - \sqrt{x_2})^2 &> 0 \end{aligned}$$

Hence, $f(x) = \sqrt{x}$ is concave and so is l_s .

- (d) Let's differentiate with respect to $y\mathbf{w}^T\mathbf{x}$. If $\text{sign}(\mathbf{w}^T\mathbf{x}) = y$, $l'_s(\mathbf{w}; \mathbf{x}, y) = 0$. If $\text{sign}(\mathbf{w}^T\mathbf{x}) \neq y$, $l'_s(\mathbf{w}; \mathbf{x}, y) = \frac{1}{2}(-y\mathbf{w}^T\mathbf{x})^{\frac{1}{2}}(-1) = \frac{1}{2\sqrt{-yy\mathbf{w}^T\mathbf{x}}}$.
To check differentiability, we need to check the limit at point 0. Let $z = y\mathbf{w}^T\mathbf{x}$. Then, $\lim_{x \rightarrow 0^-} -\frac{1}{\sqrt{z}} = -\infty$. Hence, l_s is not differentiable at $y\mathbf{w}^T\mathbf{x} = 0$.

2. Derive $\nabla l_s(w, x, y)$.

- (a) $\begin{cases} 0, & \text{if } y = \text{sign}(w^T x) \\ -\frac{yx}{2\sqrt{-y\mathbf{w}^T\mathbf{x}}}, & \text{if } y \neq \text{sign}(w^T x) \end{cases}$
- (b) $\begin{cases} 0, & \text{if } y = \text{sign}(w^T x) \\ -\frac{yx}{2\sqrt{y\mathbf{w}^T\mathbf{x}}}, & \text{if } y \neq \text{sign}(w^T x) \end{cases}$
- (c) $\begin{cases} 0, & \text{if } y = \text{sign}(w^T x) \\ \frac{yx}{2\sqrt{-y\mathbf{w}^T\mathbf{x}}}, & \text{if } y \neq \text{sign}(w^T x) \end{cases}$
- (d) $\begin{cases} 0, & \text{if } y = \text{sign}(w^T x) \\ \frac{yx}{2\sqrt{y\mathbf{w}^T\mathbf{x}}}, & \text{if } y \neq \text{sign}(w^T x) \end{cases}$

Solution:

The correct answer is (a).

Although l_s not differentiable at $y\mathbf{w}^T\mathbf{x} = 0$, the subgradient exists and hence (stochastic) gradient descent converges. To derive the subgradient let's rewrite the function l_s as $l_s(\mathbf{w}; \mathbf{x}, y) = \max(0, \sqrt{-y\mathbf{w}^T\mathbf{x}})$. Now let $f(z) = \max(0, -\sqrt{yz})$ and $g(\mathbf{w}) = \mathbf{w}^T\mathbf{x}$. We use the chain rule

$$\frac{\partial}{\partial w_i} f(g(\mathbf{w})) = \frac{\partial f}{\partial z} \frac{\partial g}{\partial w_i}$$

. We get

$$\frac{\partial f}{\partial z} = \begin{cases} 0, & \text{for } \text{sign}(z) = y \\ -\frac{y}{2\sqrt{-yz}}, & \text{for } \text{sign}(z) \neq y \end{cases}$$

and $\frac{\partial g}{\partial w_i} = x_i$. Hence,

$$\frac{\partial f(g(\mathbf{w}))}{\partial w_i} = \begin{cases} 0, & \text{for } \text{sign}(z) = y \\ -\frac{y\mathbf{x}}{2\sqrt{-y\mathbf{w}^T\mathbf{x}}}, & \text{for } \text{sign}(z) \neq y \end{cases}$$

3. The exercise suggests to train an SVM, where we penalise the margin violation given by $(1 - y\mathbf{w}^T\mathbf{x})_+ = \max(1 - y\mathbf{w}^T\mathbf{x}, 0)$, not linearly but with the square root instead. Correspondingly, our modified SVM seeks to optimise the following objective

$$L(\mathbf{w}) = \frac{1}{n} \sum_{i=1}^n \sqrt{(1 - y\mathbf{w}^T\mathbf{x})_+} + \lambda \|\mathbf{w}\|^2$$

Pick the correct update step for stochastic gradient descent.

- (a) Pick $i_t \sim \text{Unif}(1, 2, \dots, n)$.

If $y_{it}\mathbf{w}_t^T\mathbf{x}_{it} < 1$

$$w_{t+1} = w_t(1 - \eta_t 2\lambda \mathbf{w}_t) + \eta_t \frac{y_i \mathbf{x}_i}{2\sqrt{(1 - y_i \mathbf{w}_t^T \mathbf{x}_i)}}$$

Else

$$w_{t+1} = w_t(1 - \eta_t 2\lambda \mathbf{w}_t)$$

- (b) Pick $i_t \sim \text{Unif}(1, 2, \dots, n)$.

If $y_{it}\mathbf{w}_t^T\mathbf{x}_{it} < 1$

$$w_{t+1} = w_t(1 - \eta_t 2\lambda \mathbf{w}_t)$$

Else

$$w_{t+1} = w_t(1 - \eta_t 2\lambda \mathbf{w}_t) + \eta_t \frac{y_i \mathbf{x}_i}{2\sqrt{(1 - y_i \mathbf{w}_t^T \mathbf{x}_i)}}$$

- (c) Pick $i_t \sim \text{Unif}(1, 2, \dots, n)$.

If $y_{it}\mathbf{w}_t^T\mathbf{x}_{it} < 1$

$$w_{t+1} = w_t(1 + \eta_t 2\lambda \mathbf{w}_t) + \eta_t \frac{y_i \mathbf{x}_i}{2\sqrt{(1 - y_i \mathbf{w}_t^T \mathbf{x}_i)}}$$

Else

$$w_{t+1} = w_t(1 + \eta_t 2\lambda \mathbf{w}_t)$$

- (d) Pick $i_t \sim \text{Unif}(1, 2, \dots, n)$.

If $y_{it}\mathbf{w}_t^T\mathbf{x}_{it} < 1$

$$w_{t+1} = w_t(1 + \eta_t 2\lambda \mathbf{w}_t)$$

Else

$$w_{t+1} = w_t(1 + \eta_t 2\lambda \mathbf{w}_t) + \eta_t \frac{y_i \mathbf{x}_i}{2\sqrt{(1 - y_i \mathbf{w}_t^T \mathbf{x}_i)}}$$

Solution:

The correct answer is (a).

For $y_{it}\mathbf{w}_t^T\mathbf{x}_{it} < 1$,

$$\nabla_{\mathbf{w}} L = -\frac{y_i \mathbf{x}_i}{2\sqrt{(1 - y_i \mathbf{w}_t^T \mathbf{x}_i)}} + 2\lambda \mathbf{w}_t$$

Else,

$$\nabla_{\mathbf{w}} L = 2\lambda \mathbf{w}_t$$

Why may this modification not be a good idea? You can see that the weight update due to margin violations gets rescaled as a result of the modification by the factor $\frac{1}{2\sqrt{1-y_i\mathbf{w}^T\mathbf{x}_i}}$. This factor is small when the margin violation is large and large when the margin violation is small, which may make training this modified SVM troublesome.

Problem 2 (Kernels):

Use the basic rules for kernel decomposition discussed in class or otherwise and assuming that $k(x, y)$ is a valid kernel, letting $f : \mathbb{R} \rightarrow \mathbb{R}$ in a) and b), $g : \mathcal{X} \rightarrow \mathbb{R}_+$ for d), $f : \mathcal{X} \rightarrow \mathbb{R}$ for e) and f), and $\phi : \mathcal{X} \rightarrow \mathcal{X}'$.

4. Mark the following statements as True or False. Try to justify your answers to yourself.

- (a) $k_a(x, y) = f(k(x, y))$ is a valid kernel, if f is a polynomial with non-negative coefficients.
- (b) $k_b(x, y) = f(k(x, y))$ is a valid kernel, if f is any polynomial.
- (c) $k_c(x, y) = \exp(k(x, y))$ is a valid kernel.
- (d) $k_d(x, y) = g(x)k(x, y)g(y)$ is a valid kernel.
- (e) $k_e(x, y) = f(x)k(x, y)f(y)$ is a valid kernel.
- (f) $k_f(x, y) = k(\phi(x), \phi(y))$ is a valid kernel.

Solution:

(a), (c), (d), (e) and (f) are True.

- (a) Since each polynomial term is a product of kernels with non-negative coefficients, the proof follows from the rules of addition and multiplication yielding valid kernels.
- (b) Product of kernels with *negative* coefficients is not necessarily a valid kernel.
- (c) We can use the Taylor expansion around 0:

$$\begin{aligned} \exp(k(x, y)) &= \exp(0) + \exp(0)k(x, y) + \frac{\exp(0)}{2!}(k(x, y))^2 + \dots \\ &= 1 + k(x, y) + \frac{1}{2}(k(x, y))^2 + \frac{1}{6}(k(x, y))^3 \dots \end{aligned}$$

- (d) and (e) Since $k(x, y)$ is a valid kernel, we can define a feature map $\phi(\cdot)$, such that $k(x, y) = \langle \phi(x), \phi(y) \rangle$.
Now,

$$k_e(x, y) = f(x)k(x, y)f(y) = f(y)f(x)\langle \phi(x), \phi(y) \rangle = f(y)\langle f(x)\phi(x), \phi(y) \rangle = \langle f(x)\phi(x), f(y)\phi(y) \rangle$$

Hence, with the new feature map $\phi_e(\cdot) = f(\cdot)\phi(\cdot)$, $k_e(x, y)$ is a valid kernel (symmetry and positive definiteness properties don't change). This is a solution for (e). (d) follows from this, as it a specific case of the same.

- (f) We know that $k(x, y)$ is a valid kernel and hence, on any set of vectors (also transformed ones) it yields a valid kernel.

5. For $\mathbf{x}, \mathbf{x}' \in \mathbb{R}^d$, and $K(\mathbf{x}, \mathbf{x}') = (\mathbf{x}^T\mathbf{x}' + 1)^2$, identify possible feature maps $\phi(\mathbf{x})$, such that $k(\mathbf{x}, \mathbf{x}') = \phi(\mathbf{x})^T\phi(\mathbf{x}')$. Let $\mathbf{x}^T = (x_1, \dots, x_d)$.

- (a) $(1, \sqrt{2}x_1, \dots, \sqrt{2}x_d, x_1x_1, x_1x_2, \dots, x_ix_j, \dots)$
- (b) $(1 + x_1, \dots, 1 + x_i, \dots, 1 + x_d)$

- (c) $(1, -\sqrt{2}x_1, \dots, -\sqrt{2}x_d, -x_1x_1, -x_1x_2, \dots - x_ix_j \dots)$
 (d) $(1, \frac{1}{\sqrt{2}}x_1, \dots, \frac{1}{\sqrt{2}}x_d, x_1x_1, x_1x_2, \dots x_ix_j \dots, \frac{1}{\sqrt{2}}x_1, \dots, \frac{1}{\sqrt{2}}x_d)$

Solution:

(a) and (c) are correct answers.

$$(\mathbf{x}^T \mathbf{x}' + 1)^2 = (\sum_i x_i x'_i + 1)^2 = 1 + 2\sum_i x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j)$$

(a) $(1, \sqrt{2}x_1, \dots, \sqrt{2}x_d, x_1x_1, x_1x_2, \dots x_ix_j \dots)^T (1, \sqrt{2}x_1, \dots, \sqrt{2}x_d, x_1x_1, x_1x_2, \dots x_ix_j \dots)$
 $= 1 + 2\sum_i x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j)$.

(b) $(1 + x_1, \dots, 1 + x_i, \dots, 1 + x_d)^T (1 + x_1, \dots, 1 + x_i, \dots, 1 + x_d)$
 $= \sum_i (1 + x_i)^2 = \sum_i (1 + 2x_i + x_i^2) \neq 1 + 2\sum_i x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j)$.

(c) $(1, -\sqrt{2}x_1, \dots, -\sqrt{2}x_d, -x_1x_1, -x_1x_2, \dots -x_ix_j \dots)^T (1, -\sqrt{2}x_1, \dots, -\sqrt{2}x_d, -x_1x_1, -x_1x_2, \dots -x_ix_j \dots)$
 $= 1 + 2\sum_i x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j)$.

(d) $(1, \frac{1}{\sqrt{2}}x_1, \dots, \frac{1}{\sqrt{2}}x_d, x_1x_1, \dots x_ix_j \dots, \frac{1}{\sqrt{2}}x_1, \dots, \frac{1}{\sqrt{2}}x_d)^T (1, \frac{1}{\sqrt{2}}x_1, \dots, \frac{1}{\sqrt{2}}x_d, x_1x_1, \dots x_ix_j \dots, \frac{1}{\sqrt{2}}x_1, \dots, \frac{1}{\sqrt{2}}x_d)$
 $= 1 + 2\sum_i \frac{1}{2} x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j) = 1 + \sum_i x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j) \neq 1 + 2\sum_i x_i x'_i + \sum_i \sum_j (x_i x_j)(x'_i x'_j)$

6. For the dataset $X = \{\mathbf{x}_i\}_{i=1,2} = \{(-3, 4), (1, 0)\}$ and the feature map $\phi(\mathbf{x}) = [x^{(1)}, x^{(2)}, \|\mathbf{x}\|]$, calculate the Gram matrix (for a vector $\mathbf{x} \in \mathbb{R}^2$ we denote by $x^{(1)}, x^{(2)}$ its components).

(a) $\begin{pmatrix} 50 & 2 \\ 2 & 2 \end{pmatrix}$

(b) $\begin{pmatrix} 50 & 4 \\ 4 & 4 \end{pmatrix}$

(c) $\begin{pmatrix} -50 & 2 \\ 2 & 2 \end{pmatrix}$

(d) $\begin{pmatrix} 50 & 2 \\ 4 & 4 \end{pmatrix}$

Solution:

The correct answer is (a).

First, we get $\phi(x)$ for each \mathbf{x} .

(a) $\phi([-3, 4]) = (-3, 4, 5)$

(b) $\phi([1, 0]) = (1, 0, 1)$

Now we get the inner products:

(a) $\phi([-3, 4])^T \phi([-3, 4]) = 50$

(b) $\phi([-3, 4])^T \phi([1, 0]) = 2$

(c) $\phi([1, 0])^T \phi([1, 0]) = 2$

And now the Gram matrix ϕ is simply given by $\phi_{i,j} = \phi(x_i)^T \phi(x_j)$; using the above:

$$\begin{pmatrix} 50 & 2 \\ 2 & 2 \end{pmatrix}$$